




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Correspondence

Histamine release theory and roles of antihistamine in the treatment of cytokines storm of COVID-19



ARTICLE INFO

Keywords

Histamine
SARS-CoV-2
COVID-19
Cytokines storm

To the Editor,

In a recent study, authors described therapeutic options for Coronavirus Disease-19 (COVID-19) [1]. Histamine is an endogenous biogenic amine distributed ubiquitously in the cells and is present in high concentrations in the lungs, skin, and gastrointestinal tract. It acts as a local mediator in the immune system. Histamine brings about complex physiologic changes, including chemotaxis, cytokine production, and gastric acid secretion. These biologic changes occur via four G protein-coupled receptor (GPCR) subtypes: H₁ receptor (H1R), H₂ receptor (H2R), H₃ receptor (H3R), and H₄ receptor (H4R) (Table 1). H1R is expressed in various cell types, such as neurons, endothelial cells, adrenal medulla, muscle cells, hepatocytes, chondrocytes, monocytes, neutrophils, eosinophils, dendritic cells (DCs), T cells, and B cells. H1R activation leads to activation of Th1 lymphocytes, and decreased humoral immunity. H2R is expressed by parietal cells of the gastric mucosa, muscle, epithelial, endothelial, neuronal, hepatocyte, and immune cells. H2R antagonizes some of the effects mediated by H1R and leads to the relaxation of smooth muscle cells, causing vasodilation. In a murine lung inflammation model, H2R loss has an effect on invariant natural killer T (iNKT) cells, aggravating local inflammation [2].

H3R functions were identified in the central nervous system and peripheral and presynaptic receptors to control the release of histamine and other neurotransmitters. H4R is preferentially expressed in the intestine, spleen, thymus, bone marrow, peripheral hematopoietic cells, and cells of the innate and adaptive immune systems. Expression of H4R is regulated by stimulation with TNF- α , IL-6, IL-10, and IL-13, leading to inhibition of cAMP accumulation and activation of mitogen-activated protein kinases (MAPK) by H4R.

So histamine is a potent inflammatory mediator, commonly associated with allergic reactions, promoting vascular and tissue changes and possessing high chemoattractant activity. The use of selective H4R ligands and/or modulation of H1 and H4 receptor synergism may be more effective in the treatment of inflammatory conditions of the lung. Histamine also modulates the inflammatory response by acting on other cellular populations, in human lung macrophages. The binding of histamine to H1R induces production of proinflammatory cytokine IL-6 and β -glucuronidase. Blocking H4R in a model of pulmonary fibrosis alleviates the inflammatory

response, reducing Cyclooxygenase 2 (COX 2) expression and activity, leukocyte infiltration, production of Transforming growth factor beta (TGF- β) (profibrotic cytokine), and collagen deposition.

At the present, there are few studies looking into the use of antihistamine products in patients with COVID-19. In self-administered high dose oral famotidine therapy, all 10 patients had marked improvements of COVID-19 symptoms [3]. Interestingly, analysis of pharmacokinetic parameters of famotidine might indicate that it needs to be given intravenously to be effective in COVID-19 treatment given its low gastrointestinal absorption and volume of distribution [4]. In a propensity-score matched retrospective cohort study comparing famotidine cohort (84 patients) to non-famotidine cohort (1536 patients), a crude analysis showed that famotidine use was significantly associated with reduced risk for death and was independently associated with risk for death or intubation (adjusted hazard ratio (aHR) 0.42, 95% CI 0.21–0.85) [5]. The famotidine group received between 10 and 40 mg/day for a median of 5.8 days, and 72% received it orally [5]. One limitation to recognize is the risk of unmeasured confounders, particularly that sicker patients might be more likely to receive proton-pump inhibitors than H2R blockers. Although famotidine is an H2R antagonist and used mainly for peptic ulcer and gastroesophageal reflux, its potential benefit was attributed to binding and inhibiting the 3-chymotrypsin-like protease [4]. There is currently one ongoing double-blind randomized controlled trial in New York evaluating the efficacy of high dose intravenous famotidine (360 mg/day) with standard of care for a maximum of 14 days in hospitalized COVID-19 patients [6]. The H2R antagonists class also includes ranitidine cimetidine, and nizatidine. In allergic reactions, the preferred antihistamines target H1R [4]. Currently, we could not find studies evaluating the efficacy of H1R blockers in COVID-19. Histamine is a main mediator that is being released by immune system and other cells as a result of virus invasions or activation. Histamine initiates abnormal immune response leading to cytokine storm and multi-organs failure. Thus, the use of antihistaminic medications could result in a significant immune modulation which may help in the treatment of cytokine storm of COVID-19. Future studies could compare H2R antagonists with those of steroid therapy in addition to the effect of combination therapy in relation to standard therapy.

<https://doi.org/10.1016/j.tmaid.2020.101874>

Received 7 August 2020; Received in revised form 31 August 2020; Accepted 1 September 2020

Available online 3 September 2020

1477-8939/© 2020 Elsevier Ltd. All rights reserved.

Table 1
Types and functions of different histamine receptors.

	Expression in Cell Types	Function	Available studies in relation to COVID-19
Histamine 1 Receptor (H1R)	neurons, endothelial cells, adrenal medulla, muscle cells, hepatocytes, chondrocytes, monocytes, neutrophils, eosinophils, dendritic cells (DCs), T cells, and B cells	<ul style="list-style-type: none"> • activation of Th1 lymphocytes, and decreased humoral immunity 	<ul style="list-style-type: none"> • none
Histamine 2 Receptor (H2R)	parietal cells of the gastric mucosa, muscle, epithelial, endothelial, neuronal, hepatocyte, and immune cells	<ul style="list-style-type: none"> • antagonizes some of the effects mediated by H1R and leads to the relaxation of smooth muscle cells, causing vasodilation. • inhibition of CXCL10, IL-12, and TNF-α stimulation of IL-10, which is likely associated with Th2 polarization 	<ul style="list-style-type: none"> • Observational studies [3–5] • Multi-site Adaptive Trials [6]
Histamine 3 Receptor (H3R)	identified in the central nervous system and peripheral and presynaptic receptors	<ul style="list-style-type: none"> • control the release of histamine and other neurotransmitters 	<ul style="list-style-type: none"> • none
Histamine 4 Receptor (H4R)	preferentially expressed in the intestine, spleen, thymus, bone marrow, peripheral hematopoietic cells, and cells of the innate and adaptive immune systems.	<ul style="list-style-type: none"> • Activation causes chemotaxis in mast cells and eosinophils, leading to accumulation of inflammatory cells and control of cytokine secretion • increased secretion of IL-31 by Th2 cells 	<ul style="list-style-type: none"> • none

Funding source

None.

Declaration of competing interest

All authors have no conflicts of interest.

References

- [1] Jutzeler CR, Bourguignon L, Weis CV, Tong B, Wong C, Rieck B, et al. Comorbidities, clinical signs and symptoms, laboratory findings, imaging features, treatment strategies, and outcomes in adult and pediatric patients with COVID-19: a systematic review and meta-analysis. *Trav Med Infect Dis* 2020;37:101825. <https://doi.org/10.1016/j.tmaid.2020.101825>.
- [2] Ferstl R, Frei R, Barcik W, Schiavi E, Wanke K, Ziegler M, et al. Histamine receptor 2 modifies iNKT cell activity within the inflamed lung. *Allergy Eur J Allergy Clin Immunol* 2017;72. <https://doi.org/10.1111/all.13227>. 1925–35.
- [3] Casale TB, Wang J, Nowak-Wegrzyn A. Acute at home management of anaphylaxis during the Covid-19 pandemic. *J Allergy Clin Immunol Pract* 2020. <https://doi.org/10.1016/j.jaip.2020.04.022>.
- [4] Ortega JT, Serrano ML, Jastrzebska B. Class a G protein-coupled receptor antagonist famotidine as a therapeutic alternative against SARS-CoV2: an in silico analysis. *Biomolecules* 2020;10:1–19. <https://doi.org/10.3390/biom10060954>.
- [5] Freedberg DE, Conigliaro J, Wang TC, Tracey KJ, Callahan MV, Abrams JA, et al. Famotidine use is associated with improved clinical outcomes in hospitalized COVID-19 patients: a propensity score matched retrospective cohort study. *Gastroenterology* 2020. <https://doi.org/10.1053/j.gastro.2020.05.053>.
- [6] Conigliaro J. Multi-site adaptive trials for COVID-19. <https://clinicaltrials.gov/ct2/show/NCT04370262?type=Intr&cond=COVID&intr=famotidine+OR+cimetidine&draw=2&rank=1>. [Accessed 13 July 2020].

Omar Abdelhay Eldanasory

Al-Zahraa Hospital, AL-Azhar University Faculty of Medicine, Cairo, Egypt

Khalid Eljaaly

Faculty of Pharmacy, King Abdulaziz University, Jeddah, Saudi Arabia

College of Pharmacy, University of Arizona, Tucson, AZ, USA

Ziad A. Memish

King Saud Medical City, Ministry of Health, Riyadh, Saudi Arabia

Al-Faisal University, Riyadh, Saudi Arabia

Hubert Department of Global Health, Rollins School of Public Health, Emory

University, Atlanta, GA, USA

Jaffar A. Al-Tawfiq*

Specialty Internal Medicine and Quality Department, Johns Hopkins Aramco

Healthcare, Dhahran, Saudi Arabia

Indiana University School of Medicine, Indianapolis, IN, USA

Johns Hopkins University School of Medicine, Baltimore, MD, USA

* Corresponding author. Dhahran Health Center, Johns Hopkins Aramco Healthcare, P.O. Box 76, Room A-428-2, Building 61, Dhahran, 31311, Saudi Arabia.

E-mail addresses: jaffar.tawfiq@jhah.com, jaltawfi@yahoo.com (J.A. Al-Tawfiq).